


Broad-leaved Bamboo

Sasa palmata

Alien	GB	IR
2000–19	277	71
1987–99	142	13
1970–86	35	4
1930–69	12	0
pre-1930	5	0




Cut-grass

Leersia oryzoides

Native	GB	IR
2000-19	5	0
1987-99	5	0
1970-86	6	0
1930-69	10	0
pre-1930	17	0

Alien	GB	IR
2000-19	2	0
1987-99	2	0
1970-86	1	0
1930-69	2	0
pre-1930	3	0




Wood Millet

Milium effusum

Native	GB	IR
2000–19	1199	64
1987–99	1159	45
1970–86	812	19
1930–69	942	18
pre-1930	238	36


Alien	GB	IR
2000–19	8	0
1987–99	4	0
1970–86	0	0
1930–69	2	0
pre-1930	1	0



Early Millet

Milium vernale


Native	GB	IR
2000–19	●	0
1987–99	●	0
1970–86	●	0
1930–69	●	0
pre-1930	○	0



Giant Fescue

Schedonorus giganteus

Native	GB	IR
2000–19	1777	414
1987–99	1726	268
1970–86	1163	80
1930–69	1407	114
pre-1930	234	56




Wood Fescue

Drymochloa sylvatica

Native	GB	IR
2000-19	127	87
1987-99	160	62
1970-86	137	35
1930-69	93	15
pre-1930	83	30


Alien	GB	IR
2000-19	0	0
1987-99	1	0
1970-86	0	0
1930-69	1	0
pre-1930	0	0



Various-leaved Fescue

Festuca heterophylla


Alien	GB	IR
2000–19	36	3
1987–99	43	2
1970–86	35	1
1930–69	36	0
pre-1930	20	0



Rush-leaved Fescue

Festuca arenaria

Native	GB	IR
2000–19	102	1
1987–99	70	0
1970–86	55	4
1930–69	69	2
pre-1930	45	1




Viviparous Sheep's-fescue

Festuca vivipara

Native	GB	IR
2000-19	783	172
1987-99	738	153
1970-86	384	24
1930-69	609	107
pre-1930	55	22

Alien	GB	IR
2000-19	0	0
1987-99	0	0
1970-86	1	0
1930-69	0	0
pre-1930	0	0




Fine-leaved Sheep's-fescue

Festuca filiformis

Native	GB	IR
2000–19	569	44
1987–99	508	54
1970–86	336	9
1930–69	400	1
pre-1930	159	3


Alien	GB	IR
2000–19	1	0
1987–99	0	0
1970–86	0	0
1930–69	0	0
pre-1930	0	0



Huon's Fescue

Festuca huonii

Native	GB	IR
2000–19	7	0
1987–99	6	0
1970–86	5	0
1930–69	2	0
pre-1930	0	0




Blue Fescue

Festuca longifolia

Native	GB	IR
2000-19	18	0
1987-99	19	0
1970-86	11	0
1930-69	4	0
pre-1930	6	0


Alien	GB	IR
2000-19	2	0
1987-99	1	0
1970-86	0	0
1930-69	0	0
pre-1930	0	0



Hard Fescue

Festuca trachyphylla


Alien	GB	IR
2000–19	226	2
1987–99	84	0
1970–86	40	0
1930–69	52	0
pre-1930	27	0



Italian Rye-grass

Lolium multiflorum

Alien	GB	IR
2000–19	1558	369
1987–99	1488	243
1970–86	969	42
1930–69	1607	140
pre-1930	270	28




Squirreltail Fescue

Vulpia bromoides

Native	GB	IR
2000-19	1566	504
1987-99	1395	335
1970-86	832	41
1930-69	1180	286
pre-1930	292	112


Alien	GB	IR
2000-19	3	0
1987-99	2	0
1970-86	3	0
1930-69	5	0
pre-1930	7	0



Rat's-tail Fescue

Vulpia myuros

Alien	GB	IR
2000–19	1164	167
1987–99	923	48
1970–86	494	6
1930–69	554	46
pre-1930	178	55




Bearded Fescue

Vulpia ciliata

Native	GB	IR
2000–19	77	0
1987–99	85	0
1970–86	58	0
1930–69	48	0
pre-1930	33	0


Alien	GB	IR
2000–19	6	0
1987–99	8	0
1970–86	5	0
1930–69	2	0
pre-1930	2	0



Mat-grass Fescue

Vulpia unilateralis


Alien	GB	IR
2000–19	16	0
1987–99	17	0
1970–86	20	0
1930–69	24	0
pre-1930	8	0



Rough Dog's-tail

Cynosurus echinatus

Alien	GB	IR
2000–19	47	2
1987–99	45	0
1970–86	26	0
1930–69	69	3
pre-1930	97	3




Borrer's Saltmarsh-grass

Puccinellia fasciculata

Native	GB	IR
2000-19	53	9
1987-99	64	7
1970-86	50	2
1930-69	76	3
pre-1930	40	8

Alien	GB	IR
2000-19	1	0
1987-99	1	0
1970-86	5	0
1930-69	0	0
pre-1930	2	0




Stiff Saltmarsh-grass

Puccinellia rupestris

Native	GB	IR
2000-19	70	0
1987-99	76	0
1970-86	65	0
1930-69	77	0
pre-1930	90	0


Alien	GB	IR
2000-19	0	0
1987-99	0	0
1970-86	0	0
1930-69	0	0
pre-1930	4	1



Lesser Quaking-grass

Briza minor

Alien	GB	IR
2000–19	73	1
1987–99	52	0
1970–86	42	0
1930–69	40	0
pre-1930	79	0




Early Meadow-grass

Poa infirma

Native	GB	IR
2000–19	249	0
1987–99	68	0
1970–86	18	0
1930–69	18	0
pre-1930	3	0


Alien	GB	IR
2000–19	0	5
1987–99	0	1
1970–86	0	0
1930–69	0	0
pre-1930	1	0



Broad-leaved Meadow-grass

Poa chaixii


Alien	GB	IR
2000–19	98	1
1987–99	87	3
1970–86	77	1
1930–69	78	0
pre-1930	52	0



Wavy Meadow-grass

Poa flexuosa

Native	GB	IR
2000–19	6	0
1987–99	6	0
1970–86	5	0
1930–69	4	0
pre-1930	2	0




Flattened Meadow-grass

Poa compressa

Native	GB	IR
2000–19	792	0
1987–99	746	0
1970–86	501	0
1930–69	642	0
pre-1930	214	0

Alien	GB	IR
2000–19	0	26
1987–99	1	18
1970–86	0	9
1930–69	0	21
pre-1930	0	19




Glaucous Meadow-grass

Poa glauca

Native	GB	IR
2000–19	37	0
1987–99	35	0
1970–86	33	0
1930–69	45	0
pre-1930	26	0

Alien	GB	IR
2000–19	0	0
1987–99	1	0
1970–86	0	0
1930–69	0	0
pre-1930	0	0




Wood Meadow-grass

Poa nemoralis

Native	GB	IR
2000-19	1671	0
1987-99	1628	0
1970-86	1087	0
1930-69	1368	0
pre-1930	307	0

Alien	GB	IR
2000-19	1	141
1987-99	2	83
1970-86	1	36
1930-69	3	58
pre-1930	3	46




Bulbous Meadow-grass

Poa bulbosa

Native	GB	IR
2000–19	105	0
1987–99	65	0
1970–86	54	0
1930–69	39	0
pre-1930	35	0


Alien	GB	IR
2000–19	3	1
1987–99	4	1
1970–86	3	0
1930–69	4	0
pre-1930	3	0



Alpine Meadow-grass

Poa alpina


Native	GB	IR
2000–19	40	3
1987–99	46	2
1970–86	41	1
1930–69	53	1
pre-1930	36	2



Slender Cock's-foot

Dactylis polygama

Alien	GB	IR
2000–19	3	0
1987–99	1	0
1970–86	3	0
1930–69	4	0
pre-1930	0	0




Whorl-grass

Catabrosa aquatica

Native	GB	IR
2000-19	398	166
1987-99	494	177
1970-86	358	39
1930-69	563	87
pre-1930	340	116

Alien	GB	IR
2000-19	0	0
1987-99	0	1
1970-86	0	0
1930-69	0	0
pre-1930	0	1




Sea Fern-grass

Catapodium marinum

Native	GB	IR
2000–19	551	187
1987–99	387	127
1970–86	198	36
1930–69	272	61
pre-1930	90	48

Alien	GB	IR
2000–19	5	0
1987–99	7	0
1970–86	2	0
1930–69	2	0
pre-1930	1	0




Fern-grass

Catapodium rigidum

Native	GB	IR
2000-19	1196	523
1987-99	1042	375
1970-86	644	60
1930-69	849	274
pre-1930	192	94

Alien	GB	IR
2000-19	2	0
1987-99	2	0
1970-86	0	0
1930-69	0	0
pre-1930	0	0




Reed Sweet-grass

Glyceria maxima

Native	GB	IR
2000–19	1280	155
1987–99	1200	133
1970–86	837	36
1930–69	901	55
pre-1930	165	37


Alien	GB	IR
2000–19	69	7
1987–99	31	1
1970–86	11	0
1930–69	12	0
pre-1930	2	0



Floating Sweet-grass

Glyceria fluitans


Native	GB	IR
2000–19	2609	880
1987–99	2537	787
1970–86	1750	140
1930–69	2278	596
pre-1930	309	108



Hybrid Sweet-grass

Glyceria fluitans x notata

	GB	IR
2000–19	● 172	● 16
1987–99	● 328	● 45
1970–86	● 288	● 9
1930–69	● 347	● 15
pre-1930	○ 165	○ 3




Plicate Sweet-grass

Glyceria notata

Native	GB	IR
2000–19	1157	281
1987–99	1110	278
1970–86	823	52
1930–69	927	98
pre-1930	232	53


Alien	GB	IR
2000–19	2	0
1987–99	2	0
1970–86	0	0
1930–69	2	0
pre-1930	1	0



Small Sweet-grass

Glyceria declinata

Native	GB	IR
2000–19	1638	461
1987–99	1536	256
1970–86	945	45
1930–69	901	98
pre-1930	165	18




Curved Hard-grass

Parapholis incurva

Native	GB	IR
2000–19	105	1
1987–99	87	1
1970–86	57	1
1930–69	69	0
pre-1930	28	0

Alien	GB	IR
2000–19	0	0
1987–99	0	0
1970–86	1	0
1930–69	6	0
pre-1930	11	0




Wood Melick

Melica uniflora

Native	GB	IR
2000-19	1287	238
1987-99	1319	198
1970-86	960	74
1930-69	1164	61
pre-1930	228	61

Alien	GB	IR
2000-19	2	0
1987-99	1	0
1970-86	0	0
1930-69	0	0
pre-1930	2	0




Mountain Melick

Melica nutans

Native	GB	IR
2000–19	226	0
1987–99	246	0
1970–86	224	0
1930–69	226	0
pre-1930	113	0

Alien	GB	IR
2000–19	3	0
1987–99	1	0
1970–86	0	0
1930–69	1	0
pre-1930	1	0




Meadow Oat-grass

Helictochloa pratensis

Native	GB	IR
2000–19	824	0
1987–99	794	0
1970–86	593	0
1930–69	649	0
pre-1930	179	0

Alien	GB	IR
2000–19	2	0
1987–99	1	0
1970–86	1	0
1930–69	2	0
pre-1930	0	0




Downy Oat-grass

Avenula pubescens

Native	GB	IR
2000-19	1364	374
1987-99	1287	327
1970-86	914	69
1930-69	1035	144
pre-1930	280	78

Alien	GB	IR
2000-19	1	0
1987-99	0	0
1970-86	0	0
1930-69	0	0
pre-1930	0	0




French Oat-grass

Gaudinia fragilis

Native	GB	IR
2000-19	48	0
1987-99	46	0
1970-86	16	0
1930-69	6	0
pre-1930	8	0


Alien	GB	IR
2000-19	7	4
1987-99	3	4
1970-86	4	0
1930-69	4	6
pre-1930	10	5



Winter Wild-oat

Avena sterilis


Alien	GB	IR
2000–19	141	4
1987–99	174	0
1970–86	81	0
1930–69	179	0
pre-1930	49	0



Oat

Avena sativa

Alien	GB	IR
2000–19	1192	295
1987–99	699	139
1970–86	175	2
1930–69	96	3
pre-1930	55	2




Yellow Oat-grass

Trisetum flavescens

Native	GB	IR
2000–19	1546	235
1987–99	1577	246
1970–86	1060	42
1930–69	1433	178
pre-1930	217	98

Alien	GB	IR
2000–19	1	0
1987–99	2	0
1970–86	1	0
1930–69	3	0
pre-1930	4	0




Crested Hair-grass

Koeleria macrantha

Native	GB	IR
2000–19	852	230
1987–99	985	196
1970–86	614	44
1930–69	903	123
pre-1930	207	76

Alien	GB	IR
2000–19	0	0
1987–99	1	0
1970–86	1	0
1930–69	0	0
pre-1930	0	0




Somerset Hair-grass

Koeleria vallesiana

Native	GB	IR
2000–19	4	0
1987–99	4	0
1970–86	4	0
1930–69	4	0
pre-1930	2	0

Alien	GB	IR
2000–19	1	0
1987–99	1	0
1970–86	0	0
1930–69	1	0
pre-1930	0	0




Bog Hair-grass

Aristavena setacea

Native	GB	IR
2000–19	71	7
1987–99	69	8
1970–86	39	7
1930–69	63	7
pre-1930	55	7

Alien	GB	IR
2000–19	0	0
1987–99	0	0
1970–86	0	0
1930–69	0	0
pre-1930	1	0




Creeping Soft-grass

Holcus mollis

Native	GB	IR
2000–19	2510	542
1987–99	2402	453
1970–86	1539	101
1930–69	2120	340
pre-1930	249	103

Alien	GB	IR
2000–19	10	0
1987–99	7	0
1970–86	2	0
1930–69	5	0
pre-1930	1	0




Grey Hair-grass

Corynephorus canescens

Native	GB	IR
2000–19	18	0
1987–99	19	0
1970–86	18	0
1930–69	17	0
pre-1930	16	0

Alien	GB	IR
2000–19	4	0
1987–99	4	1
1970–86	3	0
1930–69	4	0
pre-1930	7	0




Silver Hair-grass

Aira caryophyllea

Native	GB	IR
2000-19	1471	516
1987-99	1494	480
1970-86	987	76
1930-69	1463	465
pre-1930	351	113

Alien	GB	IR
2000-19	1	0
1987-99	1	0
1970-86	1	0
1930-69	1	0
pre-1930	2	0




Early Hair-grass

Aira praecox

Native	GB	IR
2000–19	2147	523
1987–99	2172	494
1970–86	1215	78
1930–69	2027	428
pre-1930	288	115


Alien	GB	IR
2000–19	1	0
1987–99	0	0
1970–86	0	0
1930–69	0	0
pre-1930	0	0



Holy-grass

Hierochloa odorata


Native	GB	IR
2000–19	22	0
1987–99	15	1
1970–86	14	1
1930–69	9	1
pre-1930	4	1



Bulbous Canary-grass

Phalaris aquatica


Alien	GB	IR
2000–19	107	0
1987–99	81	0
1970–86	32	0
1930–69	12	0
pre-1930	8	0



Canary-grass

Phalaris canariensis


Alien	GB	IR
2000–19	701	71
1987–99	655	41
1970–86	421	13
1930–69	700	11
pre-1930	240	11



Lesser Canary-grass

Phalaris minor


Alien	GB	IR
2000–19	43	59
1987–99	30	5
1970–86	20	0
1930–69	37	0
pre-1930	38	0



Awned Canary-grass

Phalaris paradoxa


Alien	GB	IR
2000–19	157	1
1987–99	66	0
1970–86	30	0
1930–69	33	0
pre-1930	50	1



Black Bent

Agrostis gigantea


Alien	GB	IR
2000–19	1361	125
1987–99	1299	81
1970–86	832	14
1930–69	851	26
pre-1930	183	6



Highland Bent

Agrostis castellana


Alien	GB	IR
2000–19	87	1
1987–99	62	0
1970–86	17	0
1930–69	0	0
pre-1930	8	0



Bristle Bent

Agrostis curtisii


Native	GB	IR
2000–19	173	0
1987–99	177	0
1970–86	135	0
1930–69	173	0
pre-1930	29	0



Velvet Bent

Agrostis canina

Native	GB	IR
2000–19	1893	682
1987–99	1686	417
1970–86	724	58
1930–69	289	24
pre-1930	143	9




Brown Bent

Agrostis vinealis

Native	GB	IR
2000–19	1570	330
1987–99	1117	151
1970–86	300	10
1930–69	166	4
pre-1930	47	1

Alien	GB	IR
2000–19	1	0
1987–99	1	0
1970–86	0	0
1930–69	0	0
pre-1930	0	0




Purple Small-reed

Calamagrostis canescens

Native	GB	IR
2000–19	205	0
1987–99	162	0
1970–86	157	0
1930–69	159	0
pre-1930	120	0


Alien	GB	IR
2000–19	0	0
1987–99	0	0
1970–86	1	0
1930–69	1	0
pre-1930	1	0



Scandinavian Small-reed

Calamagrostis purpurea


Native	GB	IR
2000–19	9	2
1987–99	9	0
1970–86	7	0
1930–69	4	0
pre-1930	1	0



Narrow Small-reed

Calamagrostis stricta


Native	GB	IR
2000–19	15	1
1987–99	12	1
1970–86	15	4
1930–69	19	3
pre-1930	8	5



Scottish Small-reed

Calamagrostis scotica

Native	GB	IR
2000-19	1	0
1987-99	1	0
1970-86	1	0
1930-69	0	0
pre-1930	1	0




Nit-grass

Gastridium ventricosum

Native	GB	IR
2000–19	31	0
1987–99	33	0
1970–86	18	0
1930–69	39	0
pre-1930	137	0


Alien	GB	IR
2000–19	1	0
1987–99	3	0
1970–86	1	0
1930–69	4	0
pre-1930	15	0



Loose Silky-bent

Apera spica-venti


Alien	GB	IR
2000–19	134	2
1987–99	155	1
1970–86	141	0
1930–69	142	0
pre-1930	152	0



Dense Silky-bent

Apera interrupta

Alien	GB	IR
2000–19	41	0
1987–99	65	0
1970–86	54	0
1930–69	34	0
pre-1930	36	0




Annual Beard-grass

Polypogon monspeliensis

Native	GB	IR
2000–19	51	0
1987–99	30	0
1970–86	20	0
1930–69	18	0
pre-1930	20	0


Alien	GB	IR
2000–19	359	14
1987–99	67	0
1970–86	35	0
1930–69	44	0
pre-1930	72	5



Water Bent

Polypogon viridis

Alien	GB	IR
2000–19	865	81
1987–99	75	1
1970–86	22	0
1930–69	21	0
pre-1930	19	0




Bulbous Foxtail

Alopecurus bulbosus

Native	GB	IR
2000–19	52	0
1987–99	59	0
1970–86	44	0
1930–69	44	0
pre-1930	65	0

Alien	GB	IR
2000–19	0	0
1987–99	0	0
1970–86	1	0
1930–69	0	0
pre-1930	2	0




Orange Foxtail

Alopecurus aequalis

Native	GB	IR
2000–19	139	6
1987–99	158	5
1970–86	117	0
1930–69	150	0
pre-1930	126	0

Alien	GB	IR
2000–19	7	0
1987–99	8	0
1970–86	2	0
1930–69	2	0
pre-1930	3	0




Alpine Foxtail

Alopecurus magellanicus

Native	GB	IR
2000-19	26	0
1987-99	18	0
1970-86	21	0
1930-69	20	0
pre-1930	20	0

Alien	GB	IR
2000-19	0	0
1987-99	0	0
1970-86	0	0
1930-69	1	0
pre-1930	1	0




Smaller Cat's-tail

Phleum bertolonii

Native	GB	IR
2000–19	1664	107
1987–99	1588	44
1970–86	941	7
1930–69	1378	29
pre-1930	174	5

Alien	GB	IR
2000–19	15	0
1987–99	14	0
1970–86	6	0
1930–69	10	0
pre-1930	0	0




Alpine Cat's-tail

Phleum alpinum

Native	GB	IR
2000–19	23	0
1987–99	17	0
1970–86	19	0
1930–69	20	0
pre-1930	18	0

Alien	GB	IR
2000–19	0	0
1987–99	0	0
1970–86	0	0
1930–69	1	0
pre-1930	1	0




Purple-stem Cat's-tail

Phleum phleoides

Native	GB	IR
2000-19	11	0
1987-99	12	0
1970-86	16	0
1930-69	13	0
pre-1930	21	0

Alien	GB	IR
2000-19	0	0
1987-99	0	0
1970-86	0	0
1930-69	0	0
pre-1930	3	0




Sand Cat's-tail

Phleum arenarium

Native	GB	IR
2000–19	137	42
1987–99	149	54
1970–86	100	16
1930–69	158	32
pre-1930	99	41

Alien	GB	IR
2000–19	2	0
1987–99	0	0
1970–86	1	0
1930–69	2	0
pre-1930	3	0




Smooth Brome

Bromus racemosus

Native	GB	IR
2000–19	880	61
1987–99	663	31
1970–86	364	7
1930–69	436	27
pre-1930	269	59


Alien	GB	IR
2000–19	41	0
1987–99	15	0
1970–86	15	0
1930–69	39	0
pre-1930	80	0



Slender Soft-brome

Bromus lepidus

Alien	GB	IR
2000–19	75	5
1987–99	179	8
1970–86	166	10
1930–69	645	50
pre-1930	226	18




Hairy-brome

Bromopsis ramosa

Native	GB	IR
2000-19	1719	385
1987-99	1690	354
1970-86	1082	82
1930-69	1537	206
pre-1930	231	60

Alien	GB	IR
2000-19	1	0
1987-99	0	0
1970-86	1	0
1930-69	1	0
pre-1930	1	0




Lesser Hairy-brome

Bromopsis benekenii

Native	GB	IR
2000-19	40	0
1987-99	36	0
1970-86	26	0
1930-69	29	0
pre-1930	29	0

Alien	GB	IR
2000-19	0	0
1987-99	0	0
1970-86	0	0
1930-69	0	0
pre-1930	1	0




Upright Brome

Bromopsis erecta

Native	GB	IR
2000–19	627	40
1987–99	660	33
1970–86	456	24
1930–69	578	14
pre-1930	154	8


Alien	GB	IR
2000–19	28	0
1987–99	18	0
1970–86	20	0
1930–69	15	0
pre-1930	27	0



Hungarian Brome

Bromopsis inermis


Alien	GB	IR
2000–19	191	3
1987–99	191	0
1970–86	111	0
1930–69	79	0
pre-1930	31	0



Great Brome

Anisantha diandra


Alien	GB	IR
2000–19	793	66
1987–99	262	6
1970–86	128	2
1930–69	124	0
pre-1930	49	0



Ripgut Brome

Anisantha rigida


Alien	GB	IR
2000–19	70	2
1987–99	68	0
1970–86	39	0
1930–69	24	0
pre-1930	34	0



Compact Brome

Anisantha madritensis


Alien	GB	IR
2000–19	112	8
1987–99	61	4
1970–86	35	3
1930–69	44	2
pre-1930	63	5



California Brome

Ceratochloa carinata

Alien	GB	IR
2000–19	245	2
1987–99	169	1
1970–86	58	0
1930–69	47	0
pre-1930	19	0




Heath False-bromes

Brachypodium pinnatum

Native	GB	IR
2000-19	349	16
1987-99	482	11
1970-86	256	5
1930-69	440	9
pre-1930	96	7

Alien	GB	IR
2000-19	19	0
1987-99	30	0
1970-86	15	0
1930-69	7	0
pre-1930	7	0




Bearded Couch

Elymus caninus

Native	GB	IR
2000–19	1402	76
1987–99	1316	72
1970–86	928	41
1930–69	1125	15
pre-1930	289	26

Alien	GB	IR
2000–19	2	0
1987–99	2	0
1970–86	0	0
1930–69	0	0
pre-1930	0	0




Lyme-grass

Leymus arenarius

Native	GB	IR
2000–19	405	88
1987–99	351	60
1970–86	188	21
1930–69	261	17
pre-1930	89	18

Alien	GB	IR
2000–19	16	1
1987–99	6	0
1970–86	8	0
1930–69	14	0
pre-1930	4	0




Wood Barley

Hordelymus europaeus

Native	GB	IR
2000–19	94	1
1987–99	116	0
1970–86	108	0
1930–69	91	1
pre-1930	91	1

Alien	GB	IR
2000–19	1	0
1987–99	0	0
1970–86	0	0
1930–69	0	0
pre-1930	0	0




Sea Barley

Hordeum marinum

Native	GB	IR
2000–19	93	0
1987–99	71	0
1970–86	73	0
1930–69	75	0
pre-1930	80	0


Alien	GB	IR
2000–19	0	0
1987–99	6	0
1970–86	5	0
1930–69	4	0
pre-1930	14	0



Two-rowed Barley

Hordeum distichon s.l.


Alien	GB	IR
2000–19	1258	240
1987–99	703	82
1970–86	230	3
1930–69	75	0
pre-1930	48	2



Foxtail Barley

Hordeum jubatum


Alien	GB	IR
2000–19	153	1
1987–99	211	3
1970–86	172	2
1930–69	94	1
pre-1930	33	2



Purple Moor-grass

Molinia caerulea


Native	GB	IR
2000–19	2072	918
1987–99	2044	849
1970–86	1390	208
1930–69	1921	631
pre-1930	316	138



Bermuda-grass

Cynodon dactylon

Alien	GB	IR
2000–19	59	0
1987–99	44	0
1970–86	33	0
1930–69	48	0
pre-1930	26	0




Small Cord-grass

Spartina maritima

Native	GB	IR
2000-19	23	0
1987-99	29	0
1970-86	24	0
1930-69	39	0
pre-1930	38	0


Alien	GB	IR
2000-19	2	1
1987-99	1	0
1970-86	1	2
1930-69	2	0
pre-1930	2	0



Cockspur

Echinochloa crus-galli


Alien	GB	IR
2000–19	920	39
1987–99	238	3
1970–86	110	0
1930–69	153	1
pre-1930	83	2



Green Bristle-grass

Setaria viridis


Alien	GB	IR
2000–19	392	7
1987–99	241	32
1970–86	131	1
1930–69	155	0
pre-1930	150	1



Yellow Bristle-grass

Setaria pumila


Alien	GB	IR
2000–19	520	10
1987–99	239	2
1970–86	98	0
1930–69	60	0
pre-1930	53	0



Smooth Finger-grass

Digitaria ischaemum

Alien	GB	IR
2000–19	23	1
1987–99	9	0
1970–86	8	0
1930–69	13	0
pre-1930	14	0



Hairy Finger-grass

Digitaria sanguinalis

Alien	GB	IR
2000–19	240	1
1987–99	82	0
1970–86	57	1
1930–69	44	0
pre-1930	37	0

